



## A one shot PCV2 and M.hyo vaccine containing MetaStim<sup>®</sup>, an innovative adjuvant



- One Shot Convenience
  - Weaning is a peak stress period for pigs; a one shot vaccine helps to reduce animal stress at this crucial time
  - No mixing of vaccines required
- One Shot Innovation
  - Advanced manufacturing process<sup>1</sup>
  - Tried and trusted MetaStim<sup>®</sup> adjuvant<sup>2,3</sup>



### Dosage

- One dose of 2 ml per animal
- Intramuscularly in the neck behind the ear



### Use from 3 weeks of age



### Onset of Immunity

- 3 weeks post vaccination



### Duration of Immunity

- 23 weeks post vaccination for PCV
- 16 weeks post vaccination for M.hyo



### Withdrawal Period

- Zero days



### Pack Sizes Available

- 100 ml - 50 doses
- 250 ml - 125 doses



### Accessories

- Zoetis 2 ml Pig Injector

1. Zoetis Technical Bulletin: Innovation and Technology Contributing to the Development of Suvaxyn® Circo+MH RTU.
2. Suvaxyn MH-One SPC.

**Suvaxyn® Circo+MH RTU Emulsion for Injection for Pigs**

Contains inactivated recombinant chimeric porcine circovirus type 1 containing the porcine circovirus type 2 ORF2 protein and inactivated *Mycoplasma hyopneumoniae*, strain P-5722-3.

For active immunisation of pigs against porcine circovirus type 2 (PCV2) and *Mycoplasma hyopneumoniae*. [POM-V](#)

For information about side effects, precautions, warnings and contra-indications, please refer to the product packaging and package leaflet.

For further information please see the product's SPC or contact Zoetis UK Limited, Walton Oaks, Dorking Road, Tadworth, Surrey KT20 7NS. [www.zoetis.co.uk](http://www.zoetis.co.uk) Customer Support: 0845 300 8034

Use medicines responsibly ([www.noah.co.uk/responsible](http://www.noah.co.uk/responsible)).

Date of preparation: January 2017

AH010/17