



## Effective two shot alternative for *Mycoplasma hyopneumoniae* protection.



- Reduces frequency and severity of lung lesions
- Along with Suvaxyn MH-One, allows a flexible approach to M.hyo control (decision maker tool available)
- Aqueous based Carbopol adjuvant



### Dosage

- Two doses of 2 ml per animal
- Intramuscularly behind the ear



### 1st dose > 7 days

- 2nd dose < 10 weeks
- (2 week interval required)



### Withdrawal Period

- Zero days



### Pack Sizes Available

- 100 ml - 50 doses



### Accessories

- Zoetis 2 ml Pig Injector

**Suvaxyn® M. hyo Suspension for Injection for Pigs**

Contains inactivated *Mycoplasma hyopneumoniae*, strain P-5722-3. For the active immunisation against *Mycoplasma hyopneumoniae* infection in Feeder and Finishing pigs to reduce the frequency and severity of lung lesions. POM-V

For information about side effects, precautions, warnings and contra-indications, please refer to the product packaging and package leaflet.

For further information please see the product's SPC or contact Zoetis UK Limited, Walton Oaks, Dorking Road, Tadworth, Surrey KT20 7NS. [www.zoetis.co.uk](http://www.zoetis.co.uk) Customer Support: 0845 300 8034 Use medicines responsibly ([www.noah.co.uk/responsible](http://www.noah.co.uk/responsible)).

Date of preparation: January 2017

AH009/17