



## A single dose whole virus vaccine that stimulates effective immunity against PCV.



- A unique combination of whole virus antigen with innovative adjuvant system
- Stimulates both humoral and cellular responses – two key components of effective protection against PCV
- Reduces PCV2 viraemia for reduced signs and consequences of PCV infection
- Reduces the lost DLWG and mortality associated with PMWS



### Dosage

- One dose of 2 ml per animal
- Intramuscularly behind the ear



### Use from 21 days of age



### Onset of Immunity

- 3 weeks post vaccination



### Duration of Immunity

- 19 weeks post vaccination



### Withdrawal Period

- Zero days



### Pack Sizes Available

- 100 ml - 50 doses



### Accessories

- Zoetis 2 ml Pig Injector

**Suvaxyn® PCV Suspension for Injection for Pigs**

Contains inactivated recombinant Porcine Circovirus type 1 expressing the Porcine Circovirus type 2 ORF2 protein. For the active immunisation of pigs over the age of 3 weeks against Porcine Circovirus type 2 (PCV2) to reduce viral load in blood and lymphoid tissues, and the lesions in lymphoid tissues associated with PCV2 infection, as well as to reduce clinical signs – including loss of daily weight gain, and mortality associated with Post-Weaning Multisystemic Wasting Syndrome (PMWS). POM-V

For information about side effects, precautions, warnings and contra-indications, please refer to the product packaging and package leaflet.

For further information please see the product's SPC or contact Zoetis UK Limited, Walton Oaks, Dorking Road, Tadworth, Surrey KT20 7NS. [www.zoetis.co.uk](http://www.zoetis.co.uk) Customer Support: 0845 300 8034

Use medicines responsibly ([www.noah.co.uk/responsible](http://www.noah.co.uk/responsible)).

Date of preparation: January 2017

AH008/17