



Effective two shot alternative for *Mycoplasma hyopneumoniae* protection.



- Reduces frequency and severity of lung lesions
- Along with Suvaxyn MH-One, allows a flexible approach to M.hyo control (decision maker tool available)
- Aqueous based Carbopol adjuvant



Dosage

- Two doses of 2 ml per animal
- Intramuscularly behind the ear



1st dose > 7 days

2nd dose < 10 weeks
(2 week interval required)



Withdrawal Period

- Zero days



Pack Sizes Available

- 100 ml - 50 doses



Accessories

- Zoetis 2 ml Pig Injector

Suvaxyn® M. hyo Suspension for Injection for Pigs

Contains inactivated *Mycoplasma hyopneumoniae*, strain P-5722-3. For the active immunisation against *Mycoplasma hyopneumoniae* infection in Feeder and Finishing pigs to reduce the frequency and severity of lung lesions. POM-V

For information about side effects, precautions, warnings and contra-indications, please refer to the product packaging and package leaflet.

For further information please see the product's SPC or contact Zoetis UK Limited, Walton Oaks, Dorking Road, Tadworth, Surrey KT20 7NS. www.zoetis.co.uk Customer Support: 0845 300 8034 Use medicines responsibly (www.noah.co.uk/responsible).

Date of preparation: January 2017

AH009/17