



A one shot PCV2 and M.hyo vaccine containing MetaStim[®], an innovative adjuvant



- One Shot Convenience
 - Weaning is a peak stress period for pigs; a one shot vaccine helps to reduce animal stress at this crucial time
 - No mixing of vaccines required
- One Shot Innovation
 - Advanced manufacturing process¹
 - Tried and trusted MetaStim[®] adjuvant^{2,3}



Dosage

- One dose of 2 ml per animal
- Intramuscularly in the neck behind the ear



Use from 3 weeks of age



Onset of Immunity

- 3 weeks post vaccination



Duration of Immunity

- 23 weeks post vaccination for PCV
- 16 weeks post vaccination for M.hyo



Withdrawal Period

- Zero days



Pack Sizes Available

- 100 ml - 50 doses
- 250 ml - 125 doses



Accessories

- Zoetis 2 ml Pig Injector

1. Zoetis Technical Bulletin: Innovation and Technology Contributing to the Development of Suvaxyn® Circo+MH RTU.
2. Suvaxyn MH-One SPC.

Suvaxyn® Circo+MH RTU Emulsion for Injection for Pigs

Contains inactivated recombinant chimeric porcine circovirus type 1 containing the porcine circovirus type 2 ORF2 protein and inactivated *Mycoplasma hyopneumoniae*, strain P-5722-3.

For active immunisation of pigs against porcine circovirus type 2 (PCV2) and *Mycoplasma hyopneumoniae*. [POM-V](#)

For information about side effects, precautions, warnings and contra-indications, please refer to the product packaging and package leaflet.

For further information please see the product's SPC or contact Zoetis UK Limited, Walton Oaks, Dorking Road, Tadworth, Surrey KT20 7NS. www.zoetis.co.uk Customer Support: 0845 300 8034

Use medicines responsibly (www.noah.co.uk/responsible).

Date of preparation: January 2017

AH010/17